

TEN IDEAS FROM NONLINEAR DYNAMICS

by Stephen Omohundro
at Thinking Machines Corp.
on Oct. 31, 1984

1. MANIFOLDS
2. DEFECTS
3. STRUCTURAL STABILITY
4. FOUNDATIONS OF DYNAMICS
5. BIFURCATION THEORY
6. CHAOS
7. HAMILTONIAN SYSTEMS
8. RENORMALIZATION
9. CURVATURE
10. ADIABATIC INVARIANCE

WHERE DOES NONLINEAR DYNAMICS
ARISE IN ARTIFICIAL INTELLIGENCE?

NONLINEAR DYNAMICS
↕
NONELEPHANT ZOOLOGY

in interfacing with the world:

MACHINE MANIPULATING WORLD:

Robotics (dynamics of arms, etc.)

Speech & Sound Synthesis

Graphic Display (Hamiltonian ray tracing)

MACHINE SENSING WORLD:

Vision (early processing)

Hearing

Touch, Chemical sensing, electrodynamic, etc.

PROCESSING ALGORITHMS:

Relaxation procedures in vision
Optimization by Simulated Annealing
(chip & board layout - Kirkpatrick, Gelatt, Vecchi)
DENORAL - the test in generated test

MODELING GROSS FEATURES OF COMPLEX SYSTEMS (eg. throughput of highly parallel computer)

Statistical Mechanics

Montroll { Sears Roebuck Catalogs - log normal price
distribution w/ constant dispersion for 75 years
Freeway Traffic - laws for flow instability

Language - probability of n^{th} most
popular word is $\frac{1}{n}$ to very good
approximation $1 \leq n \leq 10,000$

Maximum Entropy

ENGINEERING OF COMPONENTS

TECHNIQUES OF DYNAMICS AS THINKING PARADIGMS (which a machine should be able to use)

FUNCTIONS AND
MAPPINGS AS OBJECTS



PROCEDURES AS
ARGUMENTS +
RETURNED VALUES

FUNCTORIAL OPERATIONS
(eg. pullback) ACT ON
MANY TYPES OF OBJECT



DATA-DRIVEN
PROGRAMMING

COORDINATE FREE
CONCEPTS
UNIVERSALITY



DATA ABSTRACTION

KEY TO ALL OF
MATHEMATICS




DELAYED EVALUATION

POINCARÉ:


To understand one instance of something, consider it in relation to nearby instances.

STATE SPACE:



one point for every state of the system


DYNAMICAL SYSTEM:



state space + dynamical vector field







LINEAR DYNAMICAL SYSTEM:

State space is a vector space:



Dynamics is a linear vector field:

eg. in 2-dimensional state spaces:



A SYSTEM IS NON-LINEAR IF ITS
DYNAMICAL VECTOR FIELD IS NON-LINEAR
OR ITS STATE SPACE IS NOT A LINEAR SPACE.

STATE SPACES ARE TYPICALLY
MANIFOLDS.

(H. Weyl ~ 1920) - a space that locally looks
like Euclidean space (i.e. $\exists$ local coordinate
patches) but globally may not.

eg Spheres: S^0 S^1 S^2 ...

Tori: T^1 T^2 ...


Multihandled tori: ...

Projective spaces:




Disc with opposite
edges identified
- like asteroids game
but with extra flip

2-dim, 2-link robot arm



space of the tip is
an annulus




interior to the
annulus, there
are 2 arm states
with a given tip
state

these 2 arm
states join
at the edge
of the annulus
to form a torus

2 links in 2-dims, both ends fixed
(think of arm state with given end effector state)

Arm



Configuration Space

•
• 2 points



• 1 point




• 1 point

3-links in 2-dims, both ends fixed




3rd link is restricted to angle - its state is a point on a closed interval



Hold 3rd link fixed $\rightarrow$ Get the 2 link problem we already understand:

3 in interior $\rightarrow$ 1+2 have 2 states

3 at endpoints $\rightarrow$ 1+2 have a single state



SO THE STATE SPACE OF
THE ARM IS A CIRCLE.

3 links, 2-dims, ends fixed

Now push the ends closer together:

When close enough - only get one state when 3 is horizontal:




Configuration space:



middle of circle collapses

Pushing the ends even further together:

3 Can't even get to the middle



Configuration space:




2 circles



4 links, 2-dims, ends fixed.




Again state of last link is an interval:



State of 1st 3 links is a point at ends, in middle we saw it was a circle:

So: state space is a 2-sphere.



4 links, 2-dim, ends fixed

Push them closer to force the 3-link circle to break in the middle:



So state space of arm is a 2-torus

2 links, 3-dims, ball joints, ends fixed




state space is a
Circle = S^1

3 links, 3-dims, ball joints ends fixed
(Feeling manifolds in your bones)



3rd joint lies in a disk
Other 2 form circles which
degenerate at edge of disk



state space is a
 S^3 -sphere = S^3

Rotate links about line joining ends to
fill up S^3 with circles - Hopf fibration



Each circle links every other one
 $\Rightarrow$ constraints on deformation
of arm.

eg Twiddling thumbs with string tied is Hopf fibration



rotor in plane - nematic liquid crystal in plane




State space is circle:



periodic parallel lines in plane - smectic

State space is torus



Arrow in 3-d fixed at 1-pt.



State space is a sphere S^2

Cigar in space - fixed at center 3-d nematic




State space is a sphere with diametrically opposite pts. identified = RP^2 2-d projective plane

can think of as disk with opposite points on edge identified:



by looking at upper hemisphere (like "asteroids" but with a flip)


Rigid body in 3-d (eg. direction of end effector)



Rotation Group

RP^3 - Real projective plane of 3-dims

Choose reference configuration - every other one is gotten by rotating about some axis. So state space is ball:




Center is reference state


Any other pt. is rotation about axis between it and center by angle = distance to center

Rotation by π & $-\pi$ give same state so ball has radius π and opposite points on surface are identified. $\Rightarrow = RP^3$

Has loops that are not smoothly contractible to a point.



Map arm into RP^3 by rotation each little slice undergoes:



If shoulder fixed & hand rotates by $2\pi \rightarrow$ arm defines non-contractible loop.

If hand rotates by 4π , can smoothly deform arm back to original state keeping hand fixed.

Sine wave on a line - 1-dim crystal



state space is a circle



lattice in plane of given orientation - 2-d crystal




state space T^2 torus



3-d crystal got T^3


Planar particle in rotationally symmetric potential:



at given radius there are
2 velocities consistent w/
energy & angular momentum
(inward & outward)

these degenerate at edge of
annulus $\rightarrow$ motion in the
phase space is on a 2 torus

Lissajou figures:




at each point there are
4 possible directions

$\rightarrow$ get 4 sheets glued
together at edges

$\rightarrow$ 2 torus

The corresponding classical systems are "billiard tables". In these 2-d examples the flow is on surfaces in phase space.

Eg. Consider $\square$. Starting w/ momentum $\nearrow$ can only get 3 others: $\nwarrow, \swarrow, \searrow$. So in phase space motion is on 4 sheets:




Bouncing off an edge means jumping sheets.



Hooking up the appropriate sheets we get a torus! The motion looks integrable - indeed $|p_x|, |p_y|$ are conserved.


Eg. Consider $\triangle$. Get 6-sheets $\rightarrow$

Again get torus $\text{torus} = \text{integrable}$.








Can show anything which tiles the plane by reflection yields a torus: So $\hexagon$ is integrable.

Eg. Circular billiards has angular mom. cons. - see constrict



WHAT HAPPENS WHEN WE TRY: $\square$, $\diamond$, $\triangle$?





These are two-holed tori. Arnold's

theorem doesn't apply since the "corners" are singularities (even though p_x^2, p_y^2 are conserved in $\square$ eg.)


Systems with flow on N -dim surfaces are called pseudo-integrable.

Put in toral billiard:  =  get 4 sheets in

phase space:    

First connect the "holes" to get skeleton:

view on the tori:




A 5-holed torus !!

Topological Defects

Spill some nematic liquid crystal on the plane:



As we describe a loop in the plane the local state describes a loop in the state space




because this loop is non-contractible, there is no way to continuously fill in the hole.

⇒ get topological pocket of higher energy
hard to get rid of since must cut a lot of bonds ⇒ local minimum of energy
defects move around & interact like particles.

In plane, defects are labeled by non-contractible loops in state space.

Push 2 together, get new one. Its loop is gotten by composing 2 original loops.

⇒ Group structure = "First Homotopy Group" $\pi_1(S)$




A.) "Defect" ideas in physics


Global structure can prevent the attainment of a desirable local structure everywhere in space. The imperfections can move around but cannot be destroyed for topological reasons.

Eg. 2-dimensional crystal



energetically desirable local structure:




what about a crystal which grows like?




we ultimately find spots like:



(path closes in perfect crystal amount by which it does not in encircling dislocation is the Burgers vector)





From a perfect crystal, one can create a pair of dislocations with oppositely pointing Burgers vectors by an internal shear:



Crystals respond to stress by moving dislocations.


Eg In 3-dimensional crystals this type of dislocation forms a line. Also get screw dislocations:



Burgers vector is parallel to dislocation line

Real crystals are made of grains spanned by a web of dislocations.

Structural properties result from motion of dislocations.






FIGURE 7B




FIGURE 7C




FIGURE 7D




FIGURE 7E

FIG. 7. Line defects in nematics. This photograph has been made by M. Kléman. Here we see clearly two kinds of line defects: the thin lines, which represent topologically stable defects (index $\pm 1/2$), and the thick lines with bright sides, which represent topologically unstable defects with a probable index ± 1 . We have defined in Ftn. 26 the Brouwer degree of a map $S_n \rightarrow S_n$: it is very intuitive in the simplest case $S_1 \xrightarrow{\Delta} S_1$, since degree (k) is the algebraic number of turns on the circle S_1 of the image $k(\varphi)$ when the azimuth φ makes one positive turn (from 0 to 2π). Similarly one defines the index of the zero $v(x_0)$ of a continuous vector field on a two-dimensional surface: in any local coordinate system in the neighborhood of x_0 , it is the degree of the correspondence h : azimuth of x (near x_0) $\rightarrow$ azimuth of $v(x)$; these two azimuths are defined modulo 2π . In the case of a field of directors $n(x)$, corresponding to the local orientation (in a plane) of a nematic, the azimuth of $n(x)$ is defined modulo π and the custom in physics is to define the index of a zero $n(x_0)$ multiples of $1/2$. Examples of index $1/2$, $-1/2$, 1 , and -1 are given in Fig. 7(b), 7(c), 7(d), and 7(e), respectively. These figures represent the field of orientation of a nematic near a line defect and is a plane perpendicular to it. The index is conserved when the perpendicular plane is moved along the line defect. It is a good

$\pi_0([G:H]) = 1$: there are no wall defects. This is the case for superfluid states and more generally for all liquid or liquid crystal phases; as a general rule singularity surfaces in liquid do not seem to be stable, as is predicted by the homotopy classification for mesomorphic phases made with chiral molecules. In these cases the isotropic liquid phase has a rotating power for optical polarization, so its symmetry group cannot contain reflections: it is $E_0(3)$, the connected Euclidian group, generated by rotations and translations. These optically active chemical compounds often have chiral mesomorphic phases (chiral smectic C or cholesterics), whose rotating power can be larger by several orders of magnitude than that of the isotropic phases. Homotopic classification would predict wall defects if non-chiral molecules [$G = E(3)$, e.g., racemics] could form chiral phases [i.e., with a global symmetry group $H < E_0(3)$]. No such phases have yet been observed: they seem to be fundamentally unstable. The situation is very different for crystals. Crystal atoms or ions have no chirality. Among the 230 crystallographic classes in three dimensions, 65 correspond to crystallographic groups $H < E_0(3)$ [among these 65 there are 11 pairs of isomorphic groups, conjugate in $\text{Aff}(3)$ but not in $\text{Aff}_0(3)$]. Crystals belonging to these 65 crystallographic classes do have topologically stable wall defects, called "twin boundaries": indeed they annihilate by pairs as required by $\pi_0([E(3):H]) = \mathbb{Z}_2$.

We define at the end of Sec. II. A "topologically stable configurations." The continuous function $\phi: V \rightarrow [G:H]$, describing the local states of the medium in the volume V , is defined everywhere on V but it is not homotopic to a constant. This can happen when the domain V occupied by the phase is not contractible—for instance, if V is the volume between two coaxial vertical cylinders, bounded by two horizontal planes. Consider this volume filled with a nematic whose molecular long axis (the director) is horizontal and everywhere orthogonal to the horizontal radial direction: hence on the boundary ∂V this axis is tangential; this is actually the case in many physical situations where the nematic molecules prefer to be parallel to the wall of the container. We have described a configuration; it is topologically stable and has no defects. If V is contractible, we have proved, in the same fashion as Eq. (29), that every continuous function $V \rightarrow X$, valued in an arbitrary topological space X , is homotopic to a constant. This would exclude the existence of topologically stable configurations if we were not imposing some boundary conditions (making V equivalent to a noncontractible space) which are naturally suggested by physics. Here we define and discuss what we shall call the "normal" configurations: those which may exist inside an ordered medium otherwise in a perfect state. Consider a (contractible) domain V occupied by an ordered phase in a perfect state everywhere outside a ball B_1 of radius a , so that the function $\phi: V \rightarrow [G:H]$ describing the local states is constant on $V - B_1$. Of course V can be extended (continuously) over B_1 with the same constant value. Given a function ϕ , constant on the boundary $\partial B_1 = S_1$ of the ball B_1 , are there any ways to extend it continuously inside B_1 which are not homotopic to the constant? Since the ball B_1 , with all points of its sur-




FIG. 8. Example of a configuration in a two-dimensional "nematic" vertically oriented outside the configuration. With the origin of coordinates at the center of the figure, the orientation at the point r makes an angle $f(r)$ with the vertical, where $f(0) = 0$ and $f(r) = \pi$ for $r \geq a$. This configuration is not topologically stable; it can be continuously transformed back into the perfect "nematic" vertically oriented. Note indeed that $[G:H] = S_1$ and $\pi_2(S_1) = 1$.

face identified, is homeomorphic to the sphere S_1 , the homotopy classes of the continuous extensions of ϕ are given by $\pi_2([G:H])$. For mesomorphic phases all the $\pi_2([G:H])$ are isomorphic to $\mathbb{Z}$ (see Table II).

Here is a classical construction of a configuration belonging to the homotopy class $n \in \mathbb{Z}$. Take as the origin of coordinates the center of B_1 , the ball of radius a . Choose a continuous function $f(t)$ such that $f(0) = 0$ and $f(t) = 1$ for $t \geq 1$; such a function can even be infinitely differentiable, e.g.,


$$f(t) = \begin{cases} 1 - e^{-t^2/(1-t^2)} & 0 \leq t \leq 1 \\ 1 & 1 \leq t \end{cases} \quad (26)$$

Then, the position of the local state at the point r is obtained from that of the local state everywhere constant in $V - B_1$ by a rotation of axis r and angle $2\pi n f(r/a)$.³⁷

As Table II shows, the values of π_2 homotopy groups predict no such configurations in superfluid ^4He but do predict them for the two superfluid phases of ^3He . These configurations may have been observed, but not the line defects, yet.

³⁷We have just defined smooth maps $S_1 \xrightarrow{f} P(R, 3) \sim \text{SO}(3)$.

The reader can easily verify, from the definition we gave of the Brouwer degree in Ftn. 26, that $\deg(g_n) = 2n$. Since S_1 is the universal, double covering of $P(R, 3)$, the homotopy class is n . We could have defined $g_n = \theta \circ h_n: S_1 \xrightarrow{h_n} \text{SU}(2) \rightarrow \text{SO}(3)$; in Appendix G we define θ and we give an explicit identification of S_1 (given by B_1 with all boundary points identified) with the elements of $\text{SU}(2)$. With this identification, h_1 is the identity map and $\deg(h_n) = n$.

<u>Space</u>	<u>Homotopy Group</u>	<u>Dislocation</u>	<u>Eg</u>
<u>1-dim</u>	π_0	point	 kinks in sine-gordon eqn.
<u>2-dim</u>	π_0	line	Grain boundary in Ising spins 
	π_1	point	 vortices, flux lines <u>Rygleigh-Reinard defects</u>
	π_2	non-removable configuration no defect	<u>smectic</u>
<u>3-dim</u>	π_0	planar	<u>crystal grain boundaries</u>
	π_1	line	<u>crystal dislocations etc.</u>
	π_2	point	<u>monopole, hedgehog in field theories</u>
	π_3	configuration	<u>Finkelstein's magnetic kink in MHD</u>
<u>4-dim</u>	$\vdots$	similar to above	
	π_3	point	<u>instantons</u>

Statistical Properties can be completely dominated by defects - they are deep local minima in energy.

Might be dangerous in stochastic search algorithms: eg. in travelling salesman



true soln



topological barrier to true soln.


Phenomena like in spin glasses with frustration.

Ex. 2-dimensional Heisenberg spin system

Model for ferromagnet

$$E = -J \sum_{\langle i,j \rangle} \cos(\phi_i - \phi_j)$$


Spin waves:




$\Rightarrow$ No spontaneous magnetization

Yet there is a phase transition!

Vortices:



Vortex-Anti-Vortex Pairs:



Kosterlitz & Thouless Picture

Vortices have same statistical mechanics as a coulomb gas of charged particles.

Model for Solid $\rightarrow$ Liquid transition!

Winfree - "The Geometry of Biological Time"

Phase singularities in: color vision, time on earth, tides, fruitfly circadian rhythm, morphogenesis, cuticle, muscle fibers, slime molds, waves in brain, eye and:

SUDDEN DEATH HEART ATTACKS:

Heart cells have phase - i.e. state space is a circle

Fibrillation = phase singularity = point on surface of heart where phase winds around 2π in encircling it:



⇒ spiral waves ⇒ BAD for owner (Mines 1914)


weak pulse shifts phase slightly, strong pulse resets the heart

Graph pulse phase vs. resulting heart phase on torus

strong pulse:



weak:



Must be pulse height and phase which is arbitrarily close to setting heart cells to anything.


Shocking heart with such a pulse is likely to set up spiral waves causing spiral waves & death.

ANDRONOV AND PONTRJAGIN (1937): STRUCTURAL STABILITY

The only physically relevant aspects of a model are those which are preserved under small perturbations.

Need to specify:

1. THE ALLOWED PERTURBATIONS
2. WHEN ARE TWO SYSTEMS EQUIVALENT?



1, 2, 3, 4 Structurally stable
5, 6, 7, 8 Codimension 1
9 Codimension 2

Related notion due to Smale (~1962):

Genericity

A property of a class of systems is generic if it is true for almost all of them.

(Almost all often means: a countable intersection of open dense sets)

In some situations the property of being structurally stable is generic.

(but not for dynamical systems on state spaces of dim 4 or greater)

EXAMPLE (Relevant to vision):

In 1955 Whitney showed there were only 3 structurally stable classes of maps from the plane to the plane. (locally)

Perturbations: to nearby smooth mappings

Equivalence: 2 maps equiv. if can smoothly change coordinates to turn one into the other.


the identity:




the fold:




the cusp:



Example: Perturb the Riemann surface for $\sqrt{}$ slightly to get 3-cusps:



Example: Perturb the "folded handkerchief" map to get a cusp and a fold:



Look at a crumpled shirt for other examples.

Clear torus:



Remove hidden line:



turn a torus:




KOENDERINK AND VAN DOORN 1982

simple model of vision: maps from smooth (2-d) surfaces of objects to 2-d surface of retina.

singularities of mapping = places where line of sight is tangent to object i.e. "edges" of object.

fly over a smooth hillside $\Rightarrow$ get "lips" event:



edge of bulge ends
at two cusps $\rightarrow$ must
bend outward


A study of artists' depictions of breasts in
nudes from the 15th century to the present
shows that the edges are always drawn bending
inward.



Figure 6. Picasso: *My model* 1945.

Artist ~~draws~~ the parabolic line on the surface.

Consider the Gauss map from a surface to the unit sphere, sending a point to the unit normal there:



Positive curvature - mapped preserving orientation

Negative curvature - mapped reversing orientation

Parabolic curves - the folds of the Gauss map

For homogeneous surfaces, the observed intensity is a function only of surface orientation.

⇒ Intensity is a smooth function on sphere.

→ Parabolic lines will generically be maxima or minima and so stand out.

Moral: Things are only structurally stable in the right class of systems


DYNAMICS

Example: Damped pendulum


$$\ddot{\theta} = \frac{g}{l} \sin \theta - \alpha \dot{\theta}$$

$$\text{or } \begin{cases} \dot{\theta} = v \\ \dot{v} = \frac{g}{l} \sin \theta - \alpha v \end{cases}$$


State space is a cylinder



Open up the cylinder to see the dynamics:



Two
fixed
points:
stable sink



unstable saddle



structural stability of the dynamical vector field:

equivalence relation: 2 v.f.'s are equivalent if $\exists$ continuous change of coordinates taking orbits of one to orbits of the other

allowed perturbations: can add any smooth vector field.

Flow box theorem (straightening out theorem):

In a neighborhood of a point where a vector field doesn't vanish, we can straighten it out by a smooth change of coordinates.
(so, away from fixed points, all v.f.'s are locally equivalent)



In 1-dim:  and  are structurally stable.


 is not.

Can see by graphing:




these intersect 0 transversally $\rightarrow$ robust under perturbation

perturbs to:




Condition for structural stability is that the derivative of the vector field at the fixed point be positive or negative, but not zero.



For higher dimensions, the eigenvalues of the linearization of the v.f. at the fixed point determine whether it is structurally stable.

Hartman's theorem: If no eigenvalues are on the imaginary axis then structurally stable and equivalent to its linearization. (denoted a hyperbolic fixed point).



of e-values on left half plane = dimension of points asymptotically attracted to f.p.

this set is called the stable manifold

of e-values on right half-plane = dimension of set of points asymptotically attracted as time goes backward.


this set is called the unstable manifold.

Stable & Unstable Manifold Thm: These sets really are manifolds, are immersed copies of Euclidean space, and are tangent to the corresponding eigenspaces.



For an attracting fixed point (all e-values on the left)
the stable manifold is called the basin of attraction.

example: damped Duffing osc.



The stable manifold of the saddle separates the
two basins of attraction and so is called a
separatrix.

Periodic Orbits:

Poincaré section:



Poincaré return map:

periodic orbit is fixed point of return map.

look at eigenvalues of linearization around fixed point.
(vector linearization)


$$(\text{map linearization}) = e$$

$\Rightarrow$ role of imaginary axis in $\mathbb{C}$ for v.s. $\Leftrightarrow$ unit circle in $\mathbb{C}$ for maps.

inside unit circle = attracting directions

outside unit circle = repelling eigendirections

on the unit circle = structurally unstable.



again define stable + unstable manifolds:

If a periodic orbit is attracting
it is called a limit cycle.


Its stable manifold is its
basin of attraction.



Bifurcation Theory

codimension 1

space of
all vector
fields →




structurally stable

1 parameter family of
systems — crosses codim 1
surface transversally →
intersection robust under
perturbations.


Eigenvalues move smoothly with parameters

⇒ fixed pt. is codim 1 if has 1 zero eigenvalue or 1 complex conj. pair




Saddle Node

or



Hopf


codimension 2




can see
codim 2
behavior in
2 parameter
families

SADDLE-NODE BIFURCATION


Eigenvalues



Phase portraits



Bifurcation Diagram



Example: intensity function of a scene - convolve with ever wider gaussians (run diffusion equation on image)




Gradient vector field bifurcates
hilltop & pass annihilate

HOPF BIFURCATION


Eigenvalues



Soft
loss of
stability



Hard
loss of
stability



Examples:

Waving Flag




River Meanders



Honey



Motor Oil



Water

CODIMENSION 2 BIFURCATIONS OF FIXED PTS.

There are five cases:

1. 2 zero evlues



Bogdanoff

2. 1 zero & 1 c.c. pair




3. 2 c.c. pairs




Not understood
(interesting stuff
occurs in 3 & 4
dimensions $\Rightarrow$ extremely
complicated)


4. Hopf changes from hard to soft:





5. 2 saddle nodes come together



describes the creation of hysteresis








Quadratic Map

$$x \rightarrow 2(cx + x^2)$$

equivalent to

logistic map
 $y \rightarrow ay(1-y)$



$$C_k = C_{\infty} + A(4.6692)^{-k}$$


Feynman
#

CODIMENSION 1 BIFURCATIONS OF PERIODIC ORBITS

A periodic orbit can be codim 1 only if 1 e-value or a c.c. pair of e-values lies on the unit circle.



saddle-node of periodic orbits



period doubling

(dripping faucet: drip-drip-drip-drip-...)

⇒ dripdrip — dripdrip — ...)


The resulting orbit itself period doubles ad infinitum and the bifurcation diagram is self-similar.




secondary Hopf bifurcation

Arnold shows that this is really codimension 2 since where you leave the unit circle is crucial.


by analogy with v.l.'s expert:



State space for 2 uncoupled limit cycles is a torus:
 dynamics is linear $\dot{\theta}_1 = \omega_1$, $\dot{\theta}_2 = \omega_2$



opened up:



If $\frac{\omega_1}{\omega_2}$ is rational, torus filled with periodic orbits: resonance





If $\frac{\omega_1}{\omega_2}$ is irrational, orbits fill torus densely:



(how quickly it fills up depends on how irrational the ratio is)


Neither of these situations is structurally stable.
 A small coupling leads to **PHASE LOCKING**.



Examples: Pendulum clocks on same wall can synchron.
 Synchronization of menstrual cycles.

irrationals lock according to their continued fraction expansion.

Secondary Hopf Bifurcation



If eigenvalues leave unit circle at $e^{2\pi i p/q}$ have a resonance of order q .


$q=1,2,3,4$ = strong resonance
does complicated things
($q \geq 4$ not completely understood)

$q \geq 5$ get phase locking in a "tongue" at every rational $\frac{p}{q}$
width $\sim$ (height) $^{(q-2)/2}$

One can remove an interval from around each rational and still have stuff left!

Take $\frac{\epsilon}{q_3}$ from $\frac{\epsilon}{q_2}$ - for ϵ small enough total removed length is small.)

$\Rightarrow$ close to bifurcation there is positive measure that is not phase locked!




Truncating continued fraction gives best possible rational estimate.

Diophantine numbers have small entries $\Leftrightarrow$ very irrational
golden mean $\frac{1}{1+\frac{1}{1+\frac{1}{1+\dots}}}$ is "most irrational"


So get curves with irrational tori between tongues
Some irrationals (Liouville numbers) phase lock immediately.
Some become Cantori.



Gap - image is smaller gap... leaves only a Cantor set.




Strong Resonance
Order 2






Strong Resonance - Order 3



Weak Resonance
Order 5 and Greater



Strong Resonance - Order 4
(there are others as well)



Classic Cantor Set

recursively remove the middle third of an interval.






giving: write # in ternary expansion
(.01211021...)

Cantor set is points with no 1's. Collapse onto an interval by changing $2 \rightarrow 1$ and viewing as binary. uncountable # of points

Measure of set is $\lim_{n \rightarrow \infty} (\frac{2}{3})^n = 0$

If you blow up set by factor of 3, get 2 copies.



Hausdorff dimension of classic cantor set = $\frac{\log 2}{\log 3} \sim .63$

Fat Cantor Set: Remove middle $\frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$

Get Cantor set with positive measure (dim 1)

(zoom in with microscope, looks denser & denser)

Undernourished Cantor Set: Remove middle $\frac{1}{3}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt[3]{3}}, \dots$
zero dimensional

(zoom in, gets thinner & thinner)

Mixture: Cut $[0, 1]$ at each $\frac{p}{q}$, separate halves by $\frac{1}{q^2}$.


positive measure

zoom in on strong irrationals, looks denser & denser

weak " , looks sparser & sparser


dimension varies wildly

CHAOS




Shaking pendulum: Sequence of clockwise, counter-clockwise can be any string of 1's & 0's with appropriate initial conditions. Knowing entire past history of $c, cc, c, c, cc, \dots$ tells nothing about next.

Consider the map: $x \rightarrow 2x \bmod 1$ of $[0, 1]$ to itself.



Or on a circle:



Call left half of interval 0, right half 1.

Claim: There is an initial condition that hops left to right according to any sequence of 0's & 1's.

So this deterministic system produces 0's and 1's as random as flipping a coin.

In binary multiplying by 2 just shifts the binary point. mod 1 lops off the integer part.

Left half of $[0, 1]$ if first digit is 0, right half if 1.

So string of 1's & 0's is digits of binary expansion

.011000101100001...

Mapping smooth dynamics to space of symbol sequences is symbolic dynamics.

$2x \bmod 1$ can't be a return map since non-invertible.

Make invertible (but not continuous) in baker's transformation:




$$(x, y) \in [0, 1] \times [0, 1]$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} 2x \bmod 1 \\ y \end{pmatrix} = \begin{cases} 2x & \text{if } x \in [0, \frac{1}{2}] \\ 2x - 1 & \text{if } x \in [\frac{1}{2}, 1] \end{cases}$$

x coordinate does $2x \bmod 1$ but don't lose information since y keeps track.

symbolic dynamics



write both x and y in binary:

$$\dots \underbrace{011000101110}_{y} . \underbrace{1011001}_{x} \dots$$

Map just shifts binary point to the right.


$\Rightarrow$ chaotic orbits of all types

$\Rightarrow$ periodic orbits of every period

$\Rightarrow \exists$ dense orbit (take $\dots 0 \underline{1} 0001 \underline{10} 11 \underline{000001} 010 \dots$)

$\Rightarrow$ periodic orbits are dense.

but not continuous.



determinant = 1


$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2x+y \\ x+y \end{pmatrix}$$

A drawing of the mapping specified by (6). After only two iterations of the mapping, one already has a crazy, mind-boggling.

Arnold's cat map

$$x \rightarrow 2x \bmod 1$$


$$y \rightarrow \begin{cases} \frac{y}{2} & \text{if } x \in [0, \frac{1}{2}) \\ \frac{y+1}{2} & \text{if } x \in [\frac{1}{2}, 1) \end{cases}$$



A drawing of the Baker's transformation.

Baker's Transformation

Smale's Horseshoe



Only interested in parts that stay in rectangle
(but could cap:)






image of red



preimage of red



Things that stay in rectangle forever = Center set $\times$ Center Set



Collapse invariant Center set
to a square to get baker's
transf.



$\Rightarrow \exists$ horseshoe $\Rightarrow \infty$ many periodic orbits of
every period, chaotic orbits

horseshoe is structurally stable


Consider a "saddle" periodic orbit



If stable & unstable manifolds intersect then transverse generically.



Transversal intersection $\Rightarrow$ horseshoe $\Rightarrow$ chaotic orbits




eg: shaking pendulum
restricted 3-body problem
asymmetric top

⋮

SHOEMAKING

How do the ∞ # of periodic orbits get formed?

→ an infinite # of saddle-nodes & period doubling cascades



→ ∞ # of attracting periodic orbits near homoclinic tangency

Unstable manifold winds around - limit is Center set



⇒ 1 homoclinic tangency means an ∞ # nearby



But stable manifold does this too:

Can get homoclinic tangency when 2 Center sets intersect

If they had positive measure and the sum of the measures was greater than the interval

⇒ CONTINUUM OF HOMOCLINK TANGENCIES


(Really they have measure 0 but can have "thickness" which implies the same thing)

Key to Chaos is stretching. Brings noise on the small to the large scale.

ex Fly & the Taffey Puller - Mixing Systems


Can make "strange attractors" - fractal dimension.

Plykin
attractor
2-d map
Axiom A






The
Solenoid


Solid Torus
↓



Genodize









$$\dot{x} = -(y+z)$$


$$\dot{y} = x + \frac{1}{3}y$$

$$\dot{z} = \frac{1}{3} + z(x-m)$$

$$m = 5.7$$

Rössler Equations






Hénon
Attractor

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} y + 1 - ax^2 \\ bx \end{pmatrix}$$

$$b = .3 \quad a = 1.4$$



b
Continuation of (a)



Lorenz
Attractor

$$\begin{aligned} \dot{x} &= -\sigma x + \sigma y \\ \dot{y} &= -xz + r x - y \\ \dot{z} &= xy - bz \end{aligned}$$

$$\sigma = 10 \quad b = 8/3 \\ r = 28$$

HAMILTONIAN SYSTEMS

Point & Rigid Body Mechanics

Ray equations of Geometric Optics

Fundamental models in every one of physics

basic model: state space is even dimensional - phase space

= configuration space variables q_i $1 \leq i \leq n$

& their conjugate momenta p_i

Poisson bracket: $\{f, g\} = \sum_{i=1}^n \left(\frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q_i} \right)$

Hamiltonian function: $H(q_1, \dots, q_n, p_1, \dots, p_n)$

Dynamics: $\dot{f} = \{f, H\}$


Special Properties:

Noether's theorem: symmetries $\leftrightarrow$ conserved quantities

$$0 = X_J \cdot H = \{H, J\} = -\{J, H\} = -\dot{J}$$

Liouville's theorem: volume preserved by flow
in fact symplectic form $\omega = \sum_i dq_i \wedge dp_i$ is

$\Rightarrow$ Eigenvalues at fixed points must look like:



No attractors! Poincaré recurrence theorem

Only solvable systems were ones with lots of symmetry.

Noether $\rightarrow$ constants of motion in involution.

If can find n constants of motion I_i st $\{I_i, I_j\} = 0$
called **INTEGRABLE**. $\Rightarrow$ can find action angle variables

$\Rightarrow$ Dynamics is on n -dim tori.

Nobody could deal w/ anything else - eg. 3-body problem


Issues clear even with 2-coupled oscillators.

Phase space is 4-dim, H cuts down to 3-dim, filled with 2-dim tori, take Poincare section.

Get elliptic  & hyperbolic  fixed points
(Must be area-preserving)

KAM theorem - for small enough perturbation of integrable system, positive measure of tori persist. In other words, frequency varies as the action does.




$\Rightarrow$ Rational tori break up into island chains



get alternating elliptic & hyperbolic points at every rational torus.

The hyperbolic pts have homoclinic behavior - horseshoes, etc.


The elliptic points are surrounded by tori that are themselves breaking - so whole picture repeated ad infinitum.






Same orbits of the standard map at $h = 0.9$

Note the intricate self-similar, "fractal" structure, where islands whose structure is a small copy of the whole, form in between separatrices which form "horseshoes" and are thus chaotic.



Hénon-Heiles system - surface of section: $V(x, y) = \frac{1}{2}(x^2 + y^2) + x^2y - \frac{y^3}{3}$
 Model for star motion in a galaxy
 Left is perturbative approximation to a "const. of motion".



Close-up of chaotic region near separatrix (dots are a single orbit)









Figure 2. Phase plots of the period-doubling conservative Feigenbaum [Eqs. (4.1)–(4.9)] that is, x , horizontally versus y , vertically. A sequence of magnifications A of the regions about the orbits of period 2^k , starts at the top, on the left, going downward. The C is the parameter in (2.6).

When the phase space is this mixture of tori and chaotic regions the system is called QUASI-INTEGRABLE




A theorem for billiards similar to K.A.M. holds only if boundary has 533 derivatives.
eg The Kitaki billiard:



whispering gallery modes - regular
quasi-integrable region

The three body problem is quasi-integrable in most regimes.




Restricted three body problem has been shown to have dynamics governed by any page torn from any book of random numbers for some initial condition.

Contour plot of an area preserving map



$p = -1.055000$ $x = -0.500000$ to -0.050000 $y = -0.055000$ to 0.055000

Stochastic map




$p = -0.4224000$ $x = -0.87$ $y = -0.73$ $z = -0.60$ $w = -0.45$ $n = 10000$

When perturbed far enough we find full-fledged chaos.

Orbits diverge exponentially from one another
→ arbitrary changes in initial conditions (eg. gravitational effect of an electron on Jupiter) can completely change subsequent evolution. Thus knowing the system's history tells us nothing about its future, though deterministic, it appears random. Eg. The Stadium billiard: 



Eg. Geodesic flow on negatively curved manifolds - Anasov.


 In the limit this becomes toral billiards with round post:  i.e. a hard-sphere gas - shown to be ergodic by Sinai.

Arnold's "cat map" (can "suspend" to give a flow)

$$\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$$

In dissipative systems find "strange attractors" →






↑
classical
trajectory

← Stadium BILLIARD eigenfunctions

RENORMALIZATION

Sometimes under the appropriate microscope, things look universal.
 Define a renormalization operator on the space of all systems.
 Study fixed pts, basins of attraction etc.


Eg. Derivative Consider the functions which vanish at zero.
 $Rf(x) = 2f(\frac{x}{2})$ Only fixed pts are linear fns.
 They are attracting.



Your derivative is which linear function you are attracted to.

Eg. Saddle node

$$R\begin{pmatrix} v_x \\ v_y \end{pmatrix}(x,y) = \begin{pmatrix} 4v_x \\ 2v_y \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$



$x = x^2, y = -y$
 codim 1 attracting
 1-dim repelling

Every curve which intersects W^s maps into W^u with enough R iterations.



So W^u is universal way to annihilate a saddle node

Eg. Feigenbaum Hump maps




for



period doubling

$$Rf(x) = \alpha f \circ f(\beta x)$$




In space of hump maps, period doubling surfaces accumulate on W^u .
 spacing is R 's unstable eigenvalue = Feigenbaum's number.

Eg. KAM and breaking of golden torus

Similar renormalization around golden mean torus yields attracting fixed pt. (KAM tori and saddle with 1-d W^u).
 Universal way to break the last KAM torus.

Greene & Mackay




no tori

only golden mean left


integrable






Some orbits of the universal map for the neighbourhood of critical white circles

Some orbits of the universal $1p^2$ at $p = -0.3$



Some orbits of the universal $1p^2$ at $p = -0.3$




Non-integrable distributions

A smooth choice of vector at each point in a space is always tangent to a smooth family of curves



There does not always exist a smooth set of surfaces tangent to a field of 2-dim (or higher) planes:

eg. in 3-dim the 2-space spanned by $\hat{u}_y$ and $\hat{u}_x + \gamma \hat{u}_z$



move in little loops
can move $\perp$ to planes


This phenomenon leads to non-holonomic constraints

- only the direction you can move in phase space is constrained
not what point you are at. (parallel parking)

Lie bracket & exterior derivative measure non-integrability.

Closely related to curvature in fiber bundles

parallel transport in
a loop - don't end up where you started



Force a crab to move the
tip of a claw in a circle
- leg doesn't return.

If model leg motion decisions locally
get holonomy - curvature in the
"fiber bundle"

SIMPLIFICATION DUE TO SEPARATION OF TIME SCALES -

Approximately:

- 1) Fast degrees of freedom feel the slow degrees as if they were constant.
- 2) Slow degrees of freedom feel only the average effect of the fast degrees.

g: Boltzmann's eqn. from BBGKY - correlation fns.

Quasilinear theory - waves & background

Molecular physics

Solid state physics

Nuclei and electrons

Dynamical System with parameter λ :

$$\dot{x} = X(x, \lambda)$$

An adiabatic invariant $I(x, \lambda)$ is a function of the state x and parameter λ whose variation over times of order



$\frac{1}{\varepsilon}$ is of order ε on solution curves

of :
$$\dot{x}(t) = X(x(t), \lambda(\varepsilon t))$$

where $\lambda(\cdot)$ is a given smooth function.

Example: SHO with slowly varying parameter

Einstein & the old quantum theory



Adiabatic invariant is the action J .

$$H = \omega J$$

$$H \sim \frac{1}{\sqrt{L}} \Rightarrow \text{force} \sim \frac{1}{L^{3/2}}$$

eg: Tuning a Guitar while a note is playing

Feel an extra force! - a pseudoforce

freq of fundamental $\sim \sqrt{\text{tension}}$

so $\frac{H}{\omega} \sim \frac{H}{\sqrt{T}} = \text{const} \Rightarrow$ potential of $\sqrt{T}$
force of $\frac{1}{\sqrt{T}}$

eg: Singing in shower as tub is filling

eg. Gyromotion:




magnetic moment is adiabatic inv.
drifts in reduced dynamics

eg. Ponderomotive force:





eg. WKB waves: (slowly varying medium)



Relation between ponderomotive force of
wave on medium and dielectric "force" of
medium on wave.

Example: Ping Pong paddle & ball



$$V \sim \frac{1}{L}$$

$$\frac{\text{momentum}}{\text{impact}} \sim V$$

$$\frac{\text{impacts}}{\text{time}} \sim \frac{V}{L}$$

$$\text{So force} \sim \frac{V^2}{L} \sim \frac{1}{L^3}$$

$$\text{potential} \sim \frac{1}{L^2}$$

This is $pV^\gamma = \text{const}$ for a 1-dim gas.

Thermodynamic potential — pseudoforces

In a formal sense the entropy is an adiabatic invariant for systems ergodic on energy surface.

Late 1800's — All potential energy is really kinetic energy of hidden degrees of freedom

van Kampen -
constrained rigid body mechanics


since Lagrange we just introduce generalised coordinates obeying constraints and solve Euler-Lagrange eqn.

Really - must be steep potential well enforcing constraint:




system oscillates rapidly in direction of constraint.

If constraint well width varies, get extra pondermotive forces in constrained motion!



Pseudoforces can stabilize unstable fixed points.


eg. shaking inverted pendulum:



if fast enough, average is:



eg. Can stabilize Rayleigh-Taylor!



eg. RF stabilization of plasma instabilities

eg. forward delta wing fighter jets



aerodynamically unstable
stabilized by fast computer